



**University of  
Zurich<sup>UZH</sup>**

**Zurich Open Repository and  
Archive**

University of Zurich  
University Library  
Strickhofstrasse 39  
CH-8057 Zurich  
[www.zora.uzh.ch](http://www.zora.uzh.ch)

---

Year: 2013

---

## **New 2H-Azirin-3-amines as Synthons for Sulfur-Heterocyclic -Amino Acids**

Räber, Joëlle L ; Stoykova, Svetlana A ; Strässler, Christof ; Heimgartner, Heinz

**Abstract:** Three new spiroheterocyclic 2H-azirin-3-amines, which contain a sulfur atom, were prepared and successfully used for the synthesis of oligopeptides containing sulfur-heterocyclic -amino acids via the 'azirine/oxazolone method'.

DOI: <https://doi.org/10.1080/10426507.2012.729114>

Posted at the Zurich Open Repository and Archive, University of Zurich

ZORA URL: <https://doi.org/10.5167/uzh-78312>

Journal Article

Accepted Version

Originally published at:

Räber, Joëlle L; Stoykova, Svetlana A; Strässler, Christof; Heimgartner, Heinz (2013). New 2H-Azirin-3-amines as Synthons for Sulfur-Heterocyclic -Amino Acids. *Phosphorus, Sulfur and Silicon and the Related Elements*, 188(4):441-445.

DOI: <https://doi.org/10.1080/10426507.2012.729114>

# New *2H*-Azirin-3-amines as Synthons for Sulfur-Heterocyclic $\alpha$ -Amino Acids

by Joëlle L. Räber, Svetlana A. Stoykova, Christoph Strässler, and Heinz Heimgartner,\*

*Institute of Organic Chemistry, University of Zürich, Winterthurerstrasse 190, CH-8057*

*Zürich, Switzerland (heimgart@oci.uzh.ch)*


## Abstract

Three new spiroheterocyclic *2H*-azirin-3-amines, which contain a sulfur atom, were prepared and successfully used for the synthesis of oligopeptides containing sulfur-heterocyclic  $\alpha$ -amino acids via the ‘azirine/oxazolone method’.

## Introduction


Since the first report on the preparation of 2,2-disubstituted *2H*-azirin-3-amines **1** by Rens and Ghosez,<sup>1</sup> these cyclic amidines have been shown to be useful synthons for  $\alpha,\alpha$ -disubstituted  $\alpha$ -amino acids in syntheses of heterocycles and peptides.<sup>2</sup> For example, the reaction with thiazolidine-2,4-dione leads to the thiadiazocine derivatives **2** via a ring enlargement reaction,<sup>3</sup> whereas the reaction with a peptide acid yields the chain-extended peptide amide **3** with a terminal 2,2-disubstituted glycine residue<sup>4</sup> (*Scheme 1*). The latter reaction was used extensively for the synthesis of conformationally restricted natural and artificial oligopeptides.<sup>5</sup>

*Scheme 1*



The so-called ‘azirine/oxazolone method’ has also been modified successfully for the synthesis of endothiopeptides of type **7** containing  $\alpha,\alpha$ -disubstituted  $\alpha$ -amino acids<sup>6</sup> (Scheme 2). Thus, thiocarboxylic acids or peptides with a terminal –COSH group react with, *e.g.*, **1a** to give thioamides **5** with an extended peptide chain. Subsequent acid-catalyzed isomerization via a 1,3-thiazole intermediate yields **6**, which can be further coupled with amino acid esters.

Scheme 2



With the aim of preparing synthons for sulfur-containing heterocyclic  $\alpha$ -amino carboxylic acids, new spiroheterocyclic 2*H*-azirin-3-amines were synthesized and used in model reactions.


## Results

The original synthesis of 2*H*-azirin-3-amines **1** starts from *N,N*-disubstituted 2,2-dialkylacetamides. The reaction with phosgene in toluene followed by addition of a base like  $Et_3N$  leads to  $\alpha$ -chloroenamines, which on treatment with  $NaN_3$  in ether yield **1**.<sup>1</sup> As in some cases the reaction of amides with phosgene was sluggish, a modified protocol via the corresponding

thioamides was developed.<sup>7</sup> Furthermore, it was shown that the selective hydrolysis of products **3** of the azirine coupling with a  $\text{-N(Me)Ph}$  group occurred already at room temperature.<sup>8</sup> Finally, a phosgene-free one-pot synthesis of **1a** was elaborated via formation of the Li-enolate of the *N*-methyl-*N*-phenyl 2,2-dialkylacetamide and subsequent treatment with diphenyl phosphorochloridate (DPPCl) and  $\text{NaN}_3$  in THF.<sup>9</sup>


With these methods we synthesized sulfur-containing spiroheterocyclic 2*H*-azirin-3-amines **1b** and **1c**, starting with tetrahydrothiophene-3-one (**8a**) and tetrahydrothiopyran-4-one (**8b**), respectively (*Scheme 3*). Transformation to the nitriles **9** by treatment with Tosmic<sup>10</sup> or in a three-step process via  $\text{NaBH}_4$  reduction, tosylation, and substitution with  $\text{NaCN}$ ,<sup>11</sup> followed by saponification and amide formation via the acid chloride<sup>10</sup> or DCC coupling,<sup>11</sup> yielded the amides **10**. The latter were used for the azirine synthesis via the Li-enolates<sup>10</sup> or, after formation of the thioamide by treatment with Lawesson reagent, via the phosgene method.<sup>11</sup>

*Scheme 3*



In an analogous manner, the sulfur-heterocyclic dipeptide synthon **1d** was prepared via the thioamide **11**<sup>12</sup> (*Scheme 4*).


*Scheme 4*



The reactivity of the new 2*H*-azirin-3-amines **1b–1d** is similar to that of earlier prepared analogues. For example, the reaction of **1b** with thiobenzoic acid in  $\text{CH}_2\text{Cl}_2$  at 0 °C led to the

corresponding S-heterocyclic N-benzoyl  $\alpha$ -aminothiocarboxamide **12** in high yield<sup>11</sup> (Scheme 5).


Scheme 5



With the aim of demonstrating the usefulness of **1b–1d** as synthons for sulfur-heterocyclic  $\alpha$ -amino acids in the preparation of sterically congested peptides, reactions with various N-protected  $\alpha$ -amino acids or peptide fragments were performed. Thus, reaction of **1b** (Tht-synthon) with Fmoc-Val-OH in  $\text{CH}_2\text{Cl}_2$  at rt gave the protected dipeptide amide **13** as a 1:1 mixture of two diastereoisomers, which could be separated by column chromatography. Then, **13** was deprotected selectively at the C- and N-terminus by treatment with 3N HCl in MeCN/ $\text{H}_2\text{O}$  and  $\text{Et}_2\text{NH}$ , respectively, leading to **14** or **15**<sup>11</sup> (Scheme 5). These dipeptides are suitable building blocks in the synthesis of oligopeptides.


Similarly, **1c** (Thp-synthon) was reacted with various N-protected  $\alpha$ -amino acids.<sup>10,13</sup> For example, coupling with Z-protected Phe-OH gave **16**, followed by selective hydrolysis of the terminal amide group (3N HCl, THF/ $\text{H}_2\text{O}$ , rt) to yield **17**. The latter was coupled with H-Val-OMe by standard methods to give **18** (Scheme 6). It was shown by X-ray crystallography and by  $^1\text{H}$ -NMR studies that these tripeptides form a  $\beta$ -turn in the crystalline state as well as in solution, stabilized by an intramolecular hydrogen bond between NH of valine and C=O of the protecting group.

Scheme 6



Finally, the reaction of **1d** (Thp-Pro-synthon) with thiobenzoic acid occurred smoothly in THF at room temperature to give the endothiodipeptide **18**, whereas the reaction with protected  $\alpha$ -amino acids such as Boc-Val-OH led to tripeptides, *e.g.* **19**<sup>12</sup> (Scheme 7). It is important to note that **19** in the crystal adopts a conformation, which is almost identical with a  $\beta$ -turn in spite of the fact that no stabilization by an intramolecular hydrogen bond is possible. This may be explained by the presence of the  $\alpha,\alpha$ -disubstituted  $\alpha$ -amino acid in analogy to the effect of aminoisobutyric acid (Aib; see ref.<sup>14</sup>).

Scheme 7



The coupling reaction of **1d** with 4-bromobenzoic acid gave **20a**, and subsequent saponification of the ester group led to the corresponding dipeptide acid, which was again reacted with **1d**, yielding tetrapeptide ester **20b**. Repetition of these two steps led to the hexapeptide **20c** in good yield (*Scheme 7*). It was shown by X-ray crystallography and  $^1\text{H}$ -NMR studies that these highly congested oligopeptides also exist in a helical conformation in the crystalline state as well as in solution.

## Conclusions

The new sulfur-containing 2*H*-azirin-3-amines can be prepared conveniently and have been shown to be useful building blocks for novel heterocyclic  $\alpha,\alpha$ -disubstituted  $\alpha$ -amino acids in the synthesis of conformationally restricted oligopeptides. On the basis of the general reactivity of 2*H*-azirin-3-amines, they may also be used for the synthesis of novel sulfur containing heterocycles.

## Acknowledgement


Financial support of our research project by the *Swiss National Science Foundation* and *F. Hoffmann-La Roche AG*, Basel, is gratefully acknowledged.

## REFERENCES


1. M. Rens and L. Ghosez, *Tetrahedron Lett.*, 3765 (1970).
2. H. Heimgartner, *Angew. Chem., Int. Ed. Engl.*, **30**, 238 (1991).
3. S. M. Ametamey and H. Heimgartner, *Helv. Chim. Acta*, **73**, 1314 (1990).

4. a) P. Wipf and H. Heimgartner, *Helv. Chim. Acta*, **71**, 140 (1988); b) P. Wipf and H. Heimgartner, *Helv. Chim. Acta*, **73**, 13 (1990).
5. a) R. T. N. Luykx, A. Linden, and H. Heimgartner, *Helv. Chim. Acta*, **86**, 4093 (2003); b) N. Pradeille, O. Zerbe, K. Möhle, A. Linden, and H. Heimgartner, *Chem. Biodivers.*, **2**, 1127 (2005); c) W. Altherr, A. Linden, and H. Heimgartner, *Chem. Biodivers.*, **4**, 1144 (2007); d) I. Dannecker-Dörig, A. Linden, and H. Heimgartner, *Helv. Chim. Acta*, **94**, 993 (2011); S. A. Stoykova, A. Linden, and H. Heimgartner, *Helv. Chim. Acta*, **95**, in press.
6. a) J. Lehmann, A. Linden, and H. Heimgartner, *Tetrahedron*, **54**, 8721 (1998); b) J. Lehmann, A. Linden, and H. Heimgartner, *Tetrahedron*, **55**, 5359 (1999); c) R. A. Breitenmoser and H. Heimgartner, *Helv. Chim. Acta*, **84**, 786 (2001); d) A. Budzowski, A. Linden, and H. Heimgartner, *Helv. Chim. Acta*, **91**, 1471 (2008); d) A. Bärtsch, B. Bischof, and H. Heimgartner, *Pol. J. Chem.*, **83**, 195 (2009).
7. a) K. Dietliker and H. Heimgartner, *Helv. Chim. Acta*, **66**, 262 (1983); b) S. Stamm and H. Heimgartner, *Helv. Chim. Acta*, **86**, 1371 (2003).
8. P. Wipf and H. Heimgartner, *Helv. Chim. Acta*, **70**, 354 (1987).
9. J. M. Villalgordo and H. Heimgartner, *Helv. Chim. Acta*, **75**, 1866 (1992).
10. C. Strässler, A. Linden, and H. Heimgartner, *Helv. Chim. Acta*, **80**, 1528 (1997).
11. J. L. Räber, K. A. Brun, H. Heimgartner, *Heterocycles*, **74**, 397 (2007).
12. S. A. Stoykova, A. Linden, H. Heimgartner, in preparation.
13. C. Strässler, A. Linden, and H. Heimgartner, unpublished results.
14. a) G. N. Ramachandran, C. Ramakrishnan, and V. Sasisekharan, *J. Mol. Biol.*, **7**, 95 (1963); b) C. Toniolo, *CRC Crit. Rev. Biochem.*, **9**, 1 (1980); c) C. Toniolo and E. Benedetti, *Trends Biochem. Sci.*, **16**, 350 (1991).







Scheme 1




Scheme 2




Scheme 3




Scheme 4



Scheme 5



Scheme 6



Scheme 7